

NEUROLOGY

Teaching Video NeuroImage: Waddling-steppage gait secondary to spinal arachnoid cyst: An exceptional surgical outcome

Pericles Maranhao-Filho, Ricardo Magalhaes Maron and Ana Lucia Zuma de Rosso

Neurology 2008;70;85-

DOI: 10.1212/01.wnl.0000312282.36135.81

This information is current as of May 12, 2008

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://www.neurology.org/cgi/content/full/70/20/e85>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2008 by AAN Enterprises, Inc. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



Teaching Video NeuroImage: Waddling-steppage gait secondary to spinal arachnoid cyst

An exceptional surgical outcome



Pericles Maranhão-Filho, MD, PhD
Ricardo Magalhães Maron, MD
Ana Lucia Zuma de Rosso, MD, PhD


Address correspondence and reprint requests to Dr. Pericles Maranhão-Filho, Department of Neurology, Hospital Universitário Clementino Fraga Filho, Av. Brigadeiro Trompowski s/n. Universidade Federal do Rio de Janeiro, RJ, Brazil 21941-590 pmaranhofilho@gmail.com

Forty-five days after a substantial physical effort, a 14-year-old boy developed low back pain and severe waddling-steppage gait (video 1). Deep tendon reflexes were abolished in the lower limbs and plantar responses inapparent. A spinal MRI depicted a cyst extending from T5 to S2 (figure). A complete thoracic cyst resection (video 2) and a lumbar cyst peritoneal shunt were performed. The pathologic specimen examination was conclusive for arachnoid cyst. After 4 months, the gait was normal (video 3). Speculatively, the physical effort accentuated a ball valve phenomenon¹ in a preexisting asymptomatic idiopathic arachnoid cyst.

REFERENCE

1. Sharma A, Sayal P, Badhe P, Pandey A, Diyora B, In-gale H. Spinal intramedullary arachnoid cyst. *Ind J Pediatr* 2004;71:e65–e67.

Figure Sagittal T1-weighted gadolinium MRI of the thoracic (A) and lumbosacral (B) spinal cord showing a large, multi-lobulated posterior subarachnoid cyst extending from T5 to S2 and displacing the spine anteriorly



Supplemental data at
www.neurology.org

From the Department of Neurology and Neurosurgery, Federal University of Rio de Janeiro, Brazil.

Disclosure: The authors report no disclosures.

Teaching Video NeuroImage: Waddling-steppage gait secondary to spinal arachnoid cyst: An exceptional surgical outcome
Pericles Maranhao-Filho, Ricardo Magalhaes Maron and Ana Lucia Zuma de Rosso
Neurology 2008;70;85-
DOI: 10.1212/01.wnl.0000312282.36135.81

This information is current as of May 12, 2008

Updated Information & Services	including high-resolution figures, can be found at: http://www.neurology.org/cgi/content/full/70/20/e85
Supplementary Material	Supplementary material can be found at: http://www.neurology.org/cgi/content/full/70/20/e85/DC1
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/misc/Permissions.shtml
Reprints	Information about ordering reprints can be found online: http://www.neurology.org/misc/reprints.shtml

